



SPECIFICATIONS



AgriBusiness®
ANIMAL NUTRITION

© Perdue AgriBusiness, LLC.

All rights reserved.
Perdue® is a registered trademark of Perdue, Inc.
6906 Zion Church Rd.
Salisbury, MD 21804

METAATEIN® is a registered trademark of Perdue, Inc.

Standard Analysis	% DM
Moisture	8.4
Dry Matter	91.6
Nitrogen	13.0
Crude Protein	81.3
Soluble Protein	24.4
Ammonia	0.0
NDIP	12.8
ADIP	12.8
Crude Fiber	1.9
ADF	2.7
NDF	7.2
Lignin	0.5
Total VFA	0.0
Lactic	0.0
Acetic	0.0
Sugar	1.2
Starch	0.6
Soluble Fiber	1.2
NFC	3.0

Protein Bioavailability	% DM
RUP % CP	89.4
RUPd % RUP	82.4
dRUP % DM	59.9
dRUP-Lys % DM	4.5
dRUP-Met % DM	0.5
dRUP-EAA % DM	31.6

Macro Mineral	% DM
Ash	5.3
Calcium	0.9
Phosphorus	0.6
Magnesium	0.1
Potassium	0.5
Sulfur	0.8
Sodium	0.3
Chloride	0.3

Trace Mineral	ppm
Iron	1,441
Manganese	9.8
Zinc	53
Copper	7.9

Fatty Acids	% DM
Fat	3.1
TFA	3.0
Glycerol	0.1
Pigment	0.1
C12:0	0.0
C14:0	0.0
C16:0	0.6
C16:1	0.1
C18:0	0.2
C18:1T	0.0
C18:1C	0.9
C18:2	1.0
C18:3	0.0
Other	0.2
Fat Type	2.0
Lipolysis Rate	154.6

Pool	% DM	Rate	Int dig
CHO A1	-	-	-
CHO A4	1.23	40.00	100.0
CHO B1	0.57	17.00	75.0
CHO B2	1.23	30.00	75.0
CHO B3	5.97	7.00	20.0
CHO C	1.20	-	-
PRO A1	-	-	-
PRO A2	24.40	1.59	100.0
PRO B1	44.13	1.05	100.0
PRO B2	-	-	-
PRO C	12.80	-	-

Amino Acids	Product		Model Specifications ¹	
	% DM	% of CP	% DM	% of CP
Arginine	3.7	4.5	3.5	4.3
Histidine	4.5	5.5	4.9	6.0
Isoleucine	1.3	1.6	1.1	1.3
Leucine	10.0	12.3	10.4	12.8
Lysine	5.7	7.0	6.1	7.5
Methionine	0.7	0.9	0.7	0.9
Phenylalanine	5.1	6.3	5.4	6.6
Threonine	3.1	3.8	3.1	3.8
Tryptophan	0.9	1.1	1.0	1.2
Valine	6.5	7.9	6.7	8.3
EAA	41.5	51.0	42.9	52.7

¹When a feed component of a dairy ration consists of a blend of feed ingredients differing in rumen undegradable protein and/or post-ruminal protein digestibility, the amino acids of the resulting mixture differ in their ruminal degradability and post ruminal digestibility. Most ration formulation software allow only a single, common set of parameters to characterize ruminal degradability and post ruminal digestibility of all essential amino acids; one cannot enter parameters specific to each amino acid. Therefore, entering the amino acid profile of the whole protein of a feed mixture results in erroneous values of its metabolizable amino acid content. To circumvent this problem, users should use the 'model specifications' amino acid values in software such as CNCPS/NASEM so that the correct supply of each metabolizable amino acid is calculated correctly.