

SPECTRUM



SPECIFICATIONS



AgriBusiness®
ANIMAL NUTRITION

© Perdue AgriBusiness, LLC.
All rights reserved.
Perdue® is a registered trademark of Perdue, Inc.
6906 Zion Church Rd.
Salisbury, MD 21804
SPECTRUM AGRIRED® is a registered trademark of Perdue, Inc.

| Standard Analysis | % DM |
|-------------------|------|
| Moisture | 6.3 |
| Dry Matter | 93.7 |
| Nitrogen | 14.6 |
| Crude Protein | 91.2 |
| Soluble Protein | 44.8 |
| Ammonia | 0.0 |
| NDIP | 6.5 |
| ADIP | 6.5 |
| Crude Fiber | 0.3 |
| ADF | 0.4 |
| NDF | 1.0 |
| Lignin | 0.1 |
| Total VFA | 0.0 |
| Lactic | 0.0 |
| Acetic | 0.0 |
| Sugar | 0.2 |
| Starch | 0.1 |
| Soluble Fiber | 0.0 |
| NFC | 0.3 |

| Protein Bioavailability | % DM |
|-------------------------|------|
| RUP % CP | 93.5 |
| RUPd % RUP | 92.4 |
| dRUP % DM | 78.8 |
| dRUP-Lys % DM | 8.3 |
| dRUP-Met % DM | 0.8 |
| dRUP-EAA % DM | 45.1 |

| Macro Mineral | % DM |
|---------------|------|
| Ash | 3.3 |
| Calcium | 0.5 |
| Phosphorus | 0.3 |
| Magnesium | 0.0 |
| Potassium | 0.3 |
| Sulfur | 0.6 |
| Sodium | 0.5 |
| Chloride | 0.9 |

| Trace Mineral | ppm |
|---------------|-------|
| Iron | 2,520 |
| Manganese | 3.6 |
| Zinc | 27 |
| Copper | 5.6 |

| Fatty Acids | % DM |
|----------------|------|
| Fat | 4.2 |
| TFA | 4.1 |
| Glycerol | 0.0 |
| Pigment | 0.0 |
| C12:0 | 0.0 |
| C14:0 | 0.0 |
| C16:0 | 0.6 |
| C16:1 | 0.0 |
| C18:0 | 2.7 |
| C18:1T | 0.0 |
| C18:1C | 0.4 |
| C18:2 | 0.2 |
| C18:3 | 0.0 |
| Other | 0.1 |
| Fat Type | 1.9 |
| Lipolysis Rate | 70.8 |

| Pool | % DM | Rate | Int dig |
|--------|-------|-------|---------|
| CHO A1 | - | - | - |
| CHO A4 | 0.17 | 40.00 | 100.0 |
| CHO B1 | 0.08 | 17.00 | 75.0 |
| CHO B2 | - | - | - |
| CHO B3 | 0.85 | 7.00 | 20.0 |
| CHO C | 0.17 | - | - |
| PRO A1 | - | - | - |
| PRO A2 | 44.82 | 0.86 | 100.0 |
| PRO B1 | 39.96 | 0.58 | 100.0 |
| PRO B2 | - | - | - |
| PRO C | 6.46 | - | - |

| Amino Acids | Product | | Model Specifications ¹ | |
|---------------|---------|---------|-----------------------------------|---------|
| | % DM | % of CP | % DM | % of CP |
| Arginine | 3.5 | 3.8 | 3.5 | 3.8 |
| Histidine | 5.7 | 6.2 | 5.8 | 6.4 |
| Isoleucine | 0.6 | 0.7 | 0.6 | 0.6 |
| Leucine | 12.1 | 13.3 | 12.4 | 13.6 |
| Lysine | 10.0 | 10.9 | 9.6 | 10.5 |
| Methionine | 0.9 | 1.0 | 0.9 | 1.0 |
| Phenylalanine | 6.4 | 7.1 | 6.6 | 7.2 |
| Threonine | 3.7 | 4.1 | 3.8 | 4.1 |
| Tryptophan | 1.1 | 1.2 | 1.1 | 1.3 |
| Valine | 7.8 | 8.5 | 7.9 | 8.7 |
| EAA | 51.9 | 56.9 | 52.2 | 57.2 |

¹When a feed component of a dairy ration consists of a blend of feed ingredients differing in rumen undegradable protein and/or post-ruminal protein digestibility, the amino acids of the resulting mixture differ in their ruminal degradability and post ruminal digestibility. Most ration formulation software allow only a single, common set of parameters to characterize ruminal degradability and post ruminal digestibility of all essential amino acids; one cannot enter parameters specific to each amino acid. Therefore, entering the amino acid profile of the whole protein of a feed mixture results in erroneous values of its metabolizable amino acid content. To circumvent this problem, users should use the 'model specifications' amino acid values in software such as CNCPS/NASEM so that the correct supply of each metabolizable amino acid is calculated correctly.